



R-CHOP 14 - Rituximab 375 / Cyclophosphamide 750 / Doxorubicin 50 / Vincristine 2 / Prednisolone 100, diffuse large B-non-Hodgkin Lymphoma, cycle 1-6

Protocol-ID: 112 V1.1 (Short), R-CHOP 14 (RITU375/CYCL750/DOXO50/VNCR2/PRED100), DLBCL, C1-6

Indication(s)

- NHL, B-Cell Type, Diffuse Large Cell; ICD-10 C83.3
- NHL, B-Cell Type, Follicular grade IIIb; ICD-10 C82.-, C82.7, C82.9

Protocol classification

- Classification: current standard
- Intensity: Standard dose
- Therapy mode: First line
- Therapy intention: curative

Cycles

Cycle length 14 days, recommended cycles: 6

Protocol sequences

- [R-CHOP 14 \(RITU375/CYCL750/DOXO50/VNCR2/PRED100\), DLBCL, C1-6 \(PID112\) -> RITU375, C7-8 \(PID1448\)](#)

Risks

- Emetogenicity (MASCC/ESMO): moderate (30-90%)
- Neutropenia: very high (>41%)
- Febrile Neutropenia: high (>20%)
- Anemia Hb below 8g/dl: high (16-30%)
- Neuropathy: CTC AE °3-4: 7%

Therapy

Hydration: Balanced Crystalloid Solution

HYD

Access: peripheral venous

Hydration before, during, or after antitumor therapy

Day	Substance	Dosage	Solution	Appl.	Inf. time	Procedure
1	Balanced Crystalloid Solution	500 ml		i.v.	60 min	60 min before Rituximab (d1)

Allergy prophylaxis: Rituximab Allergy prophylaxis (paracetamol, dimetindene)

AP

Access: peripheral venous

Day	Substance	Dosage	Solution	Appl.	Inf. time	Procedure
1	Paracetamol	1000 mg		p.o.		60 min before Rituximab (d1)
1	Dimetinden	4 mg	NaCl 0.9% 50 ml	i.v.	5 min	30 min before Rituximab (d1)

Antiemesis: Emetogenicity moderate, GRAN i.v., without DEXA d1-3

AE

Access: peripheral venous

ASCO 2015, DGHO 2016, DKG 2016, MASCC/ESMO 2016, if palonosetron not available

Day	Substance	Dosage	Solution	Appl.	Inf. time	Procedure
1	Granisetron	1 mg	NaCl 0.9% 50 ml	i.v.	5 min	15 min before Rituximab (d1) or other 5-HT3 receptor antagonist.

Supportive therapy: Mesna i.v., hour 0 (pre), p.o. 2 h, 6 h after onset Cyclophosphamide

SUP

Access: peripheral venous

Mesna 0h,2h,6h, prophylaxis of urinary tract toxicity by cyclophosphamide. At the time of oxazaphosphorin injection, 20% of the oxazaphosphorin dose is injected simultaneously as mesna. 2 and 6 h after onset, oral intake of 40% of the oxazaphosphorin dose.

Day	Substance	Dosage	Solution	Appl.	Inf. time	Procedure
1	Mesna	150 mg/m ² BSA		i.v.	1 min	1 min before Cyclophosphamide (d1)
1	Mesna	300 mg/m ² BSA		p.o.		60 min after Cyclophosphamide (d1)
1	Mesna	300 mg/m ² BSA		p.o.		5 h after Cyclophosphamide (d1)

Antineoplastic therapy: R-CHOP (RITU + CHOP d1)

CTX

Access: central venous

R-CHOP, all substances at d1

Day	Substance	Dosage	Solution	Appl.	Inf. time	Procedure
1-5	Prednisolone	100 mg		p.o.		1-0-0-0
Administer at least 60 minutes before rituximab on Day 1.						
1	Rituximab	375 mg/m ² BSA	NaCl 0.9% 500 ml	i.v.	4 h	Sequence
Init. Infusion rate 50mg/h; it can be increased by 50mg/h every 30min to max. 400mg/h. Further infusions: init. Infusion speed 100mg/h, which can be increased by 100mg/h every 30min to max. 400mg/h.						
1	Cyclophosphamide	750 mg/m ² BSA	NaCl 0.9% 500 ml	i.v.	60 min	Sequence
1	Doxorubicin	50 mg/m ² BSA	Dextrose 5% 250 ml	i.v.	15 min	Sequence
1	Vincristine	2 mg	NaCl 0.9% 50 ml	i.v.	3 min	Sequence

Hydration: Hydration after Vincristine

HYD

Access: peripheral venous

For the prevention of vein irritation.

Day	Substance	Dosage	Solution	Appl.	Inf. time	Procedure
1	Balanced Crystalloid Solution	250 ml		i.v.	15 min	0 min after Vincristine

Hematopoietic growth factors: FN risk above 20%, G-CSF long-acting, pegylated

HW

Access: - none -

Risk of febrile neutropenia (FN) >20%, ASCO 2015, DKG 2016

Day	Substance	Dosage	Solution	Appl.	Inf. time	Procedure
2	Pegfilgrastim	6 mg		subc	Bolus	24 h after Vincristine

Infection prophylaxis: Infection prophylaxis oral, lymphatic neoplasms

Access: - none -

Infection prophylaxis for continuous (weekly) administration until the end of therapy

Day	Substance	Dosage	Solution	Appl.	Inf. time	Procedure
1,3,5,8,10,12	Cotrimoxazole	960 mg		p.o.		1-0-0-0
continuous administration 3 times a week, continuing until d14 of the last cycle						
1-14	Amphotericin B	100 mg		p.o.		1-1-1-1
1 pipette of 1 ml (100mg), continue continuous administration until d14 of the last cycle.						

Substance linksLinks to substances are found [here](#).**Concomitant therapy supplements**

In contrast to the primary literature, prednisolone is used instead of prednisone because of more favorable pharmacokinetics at the same potency.

Dexamethasone for antiemesis on days 1-4 and prednisolone of allergy prophylaxis is covered by prednisolone of antitumor therapy.

The combination of an anthracycline and cyclophosphamide may be highly emetogenic in individual patients and require the addition of a neurokinin receptor antagonist. In this case, attention must be paid to the increase in plasma concentration of prednisolone and this may need to be adjusted.

Observe tumor lysis syndrome risk classification according to Cairo 2010; for LDH elevation without tumor bulk, use protocol "Tumor lysis syndrome prophylaxis, intermediate risk". In case of LDH elevation above twice the upper limit and tumor-bulk protocol use "tumor lysis syndrome prophylaxis, high risk".

Warnings

Doxorubicin: increased risk of cardiomyopathy, maximum cumulative dose 450-550 mg/m² KOF. In mediastinal irradiation, arterial hypertension for more than 5 years, age over 70 years or previous cardiac damage, maximum 400 mg/m².

For DOXO extravasation: dry cold (not just before or after Dexrazoxane infusion) on day of extravasation. Dexrazoxane i.v. for 3 days: 2 days 1000 mg/m², 3rd day 500 mg/m², do not use in parallel with DMSO. First infusion as soon as possible and within the first 6 hours.

References

- Pfreundschuh M, Six versus eight cycles of bi-weekly CHOP-14 with or without rituximab in elderly patients with aggressive CD20+ B-cell lymphomas: a randomised controlled trial (RICOVER-60). Lancet Oncol 2008 Feb;9(2):105-16. doi: 10.1016/S1470-2045(08)70002-0. PMID: 18226581. [[PMID](#)]
- Brusamolino E, Dose-dense R-CHOP-14 supported by pegfilgrastim in patients with diffuse large B-cell lymphoma: a phase II study of feasibility and toxicity. Haematologica 2006 Apr;91(4):496-502. PMID: 16537117. [[PMID](#)]
- Cunningham D, Rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisolone in patients with newly diagnosed diffuse large B-cell non-Hodgkin lymphoma: a phase 3 comparison of dose intensification with 14-day versus 21-day cycles. Lancet 2013 May 25;381(9880):1817-26. doi: 10.1016/S0140-6736(13)60313-X. PMID: 23615461. [[PMID](#)]

Recommendations

- 10/2015: [European Society for Medical Oncology](#)
- 05/2021: [National Comprehensive Cancer Network](#)



The publishers and authors assume no liability for the accuracy of the contents. The application is at the own responsibility of the treating physician. ©Onkopti.

Valid since: 19.01.2015