



Atezolizumab 1200 / Carboplatin 6 / Nab-Paclitaxel 100, Non-Small Cell Lung Cancer

Protocol-ID: 1238 V1.2 (Short), ATEZ1200/CRBP6/NPAC100, NSCLC

Indication(s)

- Lung Carcinoma, Non-Small Cell (non-squamous); ICD-10 C34.-

Protocol classification

- Classification: alternative
- Intensity: Standard dose
- Therapy mode: First line
- Therapy intention: palliative

Cycles

Cycle length 21 days, recommended cycles: 6

Protocol sequences

- [IMpower130: ATEZ1200/CRBP6/NPAC100, NSCLC \(PID1238\) -|- ATEZ1200/CRBP6/NPAC100 - ATEZ1200 maint. \(PID1272\)](#)

Risks

- Emetogenicity (MASCC/ESMO): low (10-30%) Nab-paclitaxel
- Emetogenicity (MASCC/ESMO): high (>90%) Carboplatin combination
- Neutropenia: very high (>41%) °3-4: 44%
- Thrombocytopenia below 50 000/μl: very high (>41%) °3-4: 45%
- Anemia Hb below 8g/dl: high (16-30%) °3-4: 29%
- Diarrhea: CTC AE °3-4: 5%
- Fatigue: CTC AE °3-4: 6%
- Nausea: CTC AE °3-4: 3%

Therapy

Hydration: Balanced Crystalloid Solution

HYD

Access: peripheral venous

Hydration before, during, or after antitumor therapy

Day	Substance	Dosage	Solution	Appl.	Inf. time	Procedure
1	Balanced Crystalloid Solution	500 ml		i.v.	60 min	60 min before Atezolizumab (d1)
8,15	Balanced Crystalloid Solution	500 ml		i.v.	60 min	60 min before Nab-paclitaxel (d8,15)

Antiemesis: Emetogenicity high (CRBP), FOSAP, GRAN i.v., DEXA i.v.**AE**

Access: peripheral venous

DGHO 2016, DKG 2016, MASCC/ESMO 2016, carboplatin-containing combination therapies

Day	Substance	Dosage	Solution	Appl.	Inf. time	Procedure
1	Fosaprepitant	150 mg	NaCl 0.9% 150 ml	i.v.	20 min	30 min before Atezolizumab (d1)
1	Dexamethasone	12 mg	NaCl 0.9% 50 ml	i.v.	5 min	30 min before Atezolizumab (d1)
1	Granisetron	1 mg	NaCl 0.9% 50 ml	i.v.	5 min	15 min before Atezolizumab (d1)
8,15	Granisetron	1 mg	NaCl 0.9% 50 ml	i.v.	5 min	15 min before Nab-paclitaxel (d8,15)

Antineoplastic therapy: ATEZ1200/CRBP6/NPAC100**CTX**

Access: peripheral venous

Atezolizumab, carboplatin, and nab-paclitaxel in non-small cell, non-plate epithelial lung cancer

Day	Substance	Dosage	Solution	Appl.	Inf. time	Procedure
1	Atezolizumab	1200 mg	NaCl 0.9% 250 ml	i.v.	60 min	Sequence
If the first infusion was well tolerated, the second infusion can be given over 30 minutes.						
1	Carboplatin	6 AUC	Dextrose 5% 250 ml	i.v.	30 min	Sequence
1,8,15	Nab-paclitaxel	100 mg/m ² BSA	none	i.v.	30 min	Sequence

Concomitant therapy supplements

For highly emetogenic chemotherapy, additional olanzapine is recommended in the acute (day 1) and delayed phases (days 2-4) at a dosing of 5-10 mg per day (NCCN, ESMO, ASCO, Onkopedia; as of 6/24).

Granisetron instead of Dexamethasone for antiemesis on days 8 and 15 to avoid immunosuppression and the risk of infection due to Dexamethasone exposure.

References

- West H, Atezolizumab in combination with carboplatin plus nab-paclitaxel chemotherapy compared with chemotherapy alone as first-line treatment for metastatic non-squamous non-small-cell lung cancer (IMpower130): a multicentre, randomised, open-label, phase 3 trial., *Lancet Oncol* 2019 Jul;20(7):924-937. doi: 10.1016/S1470-2045(19)30167-6. PMID: 31122901. [PMID]
- Arbour KC, Impact of Baseline Steroids on Efficacy of Programmed Cell Death-1 and Programmed Death-Ligand 1 Blockade in Patients With Non-Small-Cell Lung Cancer., *J Clin Oncol* 2018 Oct 01;36(28):2872-2878. doi: 10.1200/JCO.2018.79.0006. PMID: 30125216. [PMID]

Recommendations

- 01/2023: [European Society for Medical Oncology](#)
- 02/2024: [National Comprehensive Cancer Network](#)



The publishers and authors assume no liability for the accuracy of the contents. The application is at the own responsibility of the treating physician. ©Onkopti.

Valid since: 15.02.2024