



## FOLFIRINOX - Oxaliplatin 85 / Folinic Acid 400 / Irinotecan 180 / Fluorouracil 2400, Pancreatic Cancer

Protocol-ID: 47 V1.4 (Basic), FOLFIRINOX (OXAL85/CFOL400/IRIN180/FU2400), Pankreas-Ca

### Indication(s)

- Pancreatic Cancer; ICD-10 C25.-

### Cycles

Cycle length 14 days, recommended cycles: 12

### References

- Conroy T, FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer.; N Engl J Med 2011 May 12;364(19):1817-25. doi: 10.1056/NEJMoa1011923. PMID: 21561347. [\[PMID\]](#)
- Conroy T, Irinotecan Plus Oxaliplatin and Leucovorin-Modulated Fluorouracil in Advanced Pancreatic Cancer, A Groupe Tumeurs Digestives of the Fédération Nationale des Centres de Lutte Contre le Cancer Study; J Clin Oncol 2005 Feb 20;23(6):1228-36. doi: 10.1200/JCO.2005.06.050. PMID: 15718320. [\[PMID\]](#)
- Ducreux M, Cancer of the pancreas: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann Oncol 2015 Sep;26 Suppl 5:v56-68. doi: 10.1093/annonc/mdv295. PMID: 26314780. [\[PMID\]](#)

### Recommendations

- 11/2023: [European Society for Medical Oncology](#)
- 06/2023: [National Comprehensive Cancer Network](#)

### Important notice

The copyrighted protocols are treatment recommendations. The information contained in this compilation on cytostatic drugs, concomitant medication and other therapeutic procedures, as well as dosage and application information, is continuously reviewed with all due care by the authors and editors involved. Nevertheless, the publishers and authors do not assume any liability for the correctness - also with regard to possible printing errors.

The protocols may not be changed in terms of content. Any further use of the protocols in physical or non-physical form, such as copying, distribution, disclosure, export to other media, or publication, even in excerpt form, is not permitted.

Diagnosis, indication for therapy and treatment of malignant diseases must be carried out in each individual case by the hematologist and oncologist on his or her own responsibility. The treating physician is obligated to this personal responsibility to weigh in each case before a diagnostic or therapeutic measure, indication, contraindications, dosage and application under consideration of the specialized information or other documents of the manufacturers. This applies in particular to rarely used preparations or preparations that are new to the market.



The publishers and authors assume no liability for the accuracy of the contents. The application is at the own responsibility of the treating physician. ©Onkopti.

Valid since: 20.09.2024